

Arizona Basin Outlook Report February 1, 2004



Basin Outlook Reports

and

Federal - State - Private

Cooperative Snow Surveys

For more water supply and resource management information, contact:

Larry P. Martinez

Water Supply Specialist

3003 N. Central Ave., Suite 800

Phoenix, AZ 85012-2945

(602) 280-8841

Email: Larry.Martinez@az.usda.gov

How forecasts are made

Most of the annual streamflow in the western United States originates as snowfall that has accumulated in the mountains during the winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Measurements of snow water equivalent at selected manual snow courses and automated SNOTEL sites, along with precipitation and streamflow values are used in computerized statistical and simulation models to prepare runoff forecasts. These forecasts are coordinated between hydrologists in the Natural Resources Conservation Service and the National Weather Service. Unless otherwise specified, all forecasts are for flows that would occur naturally without any upstream influences.

Forecasts of any kind, of course, are not perfect. Streamflow forecast uncertainty arises from three primary sources: (1) uncertain knowledge of future weather conditions, (2) uncertainty in the forecasting procedure, and (3) errors in the data. The forecast, therefore, must be interpreted not as a single value but rather as a range of values with specific probabilities of occurrence. The middle of the range is expressed by the 50% exceedance probability forecast, for which there is a 50% chance that the actual flow will be above, and a 50% chance that the actual flow will be below, this value. To describe the expected range around this 50% value, four other forecasts are provided, two smaller values (90% and 70% exceedance probability) and two larger values (30%, and 10% exceedance probability). For example, there is a 90% chance that the actual flow will be more than the 90% exceedance probability forecast. The others can be interpreted similarly.

The wider the spread among these values, the more uncertain the forecast. As the season progresses, forecasts become more accurate, primarily because a greater portion of the future weather conditions become known; this is reflected by a narrowing of the range around the 50% exceedance probability forecast. Users should take this uncertainty into consideration when making operational decisions by selecting forecasts corresponding to the level of risk they are willing to assume about the amount of water to be expected. If users anticipate receiving a lesser supply of water, or if they wish to increase their chances of having an adequate supply of water for their operations, they may want to base their decisions on the 90% or 70% exceedance probability forecasts, or something in between. On the other hand, if users are concerned about receiving too much water (for example, threat of flooding), they may want to base their decisions on the 30% or 10% exceedance probability forecasts, or something in between. Regardless of the forecast value users choose for operations, they should be prepared to deal with either more or less water. (Users should remember that even if the 90% exceedance probability forecast is used, there is still a 10% chance of receiving less than this amount.) By using the exceedance probability information, users can easily determine the chances of receiving more or less water.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C., 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

Issued by

Bruce I. Knight
Chief
Natural Resources Conservation Service
U.S. Department of Agriculture

Released by

Michael Somerville
State Conservationist
Natural Resources Conservation Service
Phoenix, Arizona

Arizona
Basin Outlook Report
Natural Resources Conservation Service
Phoenix, AZ

ARIZONA

Water Supply Outlook Report as of February 1, 2004

A full range of Snow Survey and Water Supply Forecasting products is available on the Arizona NRCS Home Page

Snow Survey Program

<http://www.az.nrcs.usda.gov/snow/index.html>

Helpful Internet Sites

Defending Against Drought – NRCS

<http://www.nrcs.usda.gov/feature/highlights/drought.html>

- Ideas on water, land, and crop management for you to consider while creating your drought plan.

Arizona Agri-Weekly

<http://www.nass.usda.gov/az/cur-agwk.pdf>

- Provides an overview of Arizona's crop, livestock, range and pasture conditions as reported by local staffs of the USDA's Agricultural Statistic Service and University of Arizona.

SUMMARY

Snow measurements confirm that Arizona's snowpack levels have increased since Jan. 1. Statewide, the snowpack is 63% of the 30-yr. average, compared with 41% on Jan. 1. While overall snowpack numbers have increased, it is important to note that all basins tracked in this report lag behind the 30-yr. average for Feb. 1. As a result, water managers can expect short water supplies this season.


SNOWPACK

Key Watersheds	Percent (%) of 30-Yr. Average Snowpack Levels as of February 1
Salt River Basin	62%
Verde River Basin	50%
Little Colorado River Basin	55%
San Francisco-Upper Gila River Basin	60%
Other Points of Interest	
Chuska Mountains	73%
Central Mogollon Rim	57%
Grand Canyon	73%
San Francisco Peaks	69%
Statewide Snowpack	63%

PRECIPITATION

Mountain data, from 15 NRCS SNOTEL sites, show that precipitation catch for January is below the 30-yr. average at all sites. For more information, please refer to the precipitation bar graphs found in this report.

RESERVOIR



Key storage volumes displayed in thousands of acre-feet(1000x) :

RESERVOIR	CURRENT STORAGE	LAST YEAR STORAGE	30-YEAR AVERAGE
-----	-----	-----	-----
Salt River System	842.5	558.8	1189.3
Verde River System	117.0	62.7	150.8
San Carlos Reservoir	27.2	36.1	421.8
Lyman Lake	2.1	2.2	14.7
Show Low Lake	3.2	2.0	2.9
Lake Pleasant	552.1	452.9	----
Lake Havasu	510.6	537.1	551.8
Lake Mohave	1623.4	1705.3	1672.3
Lake Mead	15434.0	16854.0	21992.0
Lake Powell	10984.0	13269.0	18463.0

STREAMFLOW



The long-term forecast (Feb-May) calls for well below normal runoff this season due to poor snowpack levels in key watersheds.



SALT RIVER BASIN as of February 1, 2004

Well below median streamflow levels are forecast for the basin. In the Salt River, near Roosevelt, the forecast calls for 54 % of median streamflow levels through MAY, while in Tonto Creek, the forecast calls for 30 % of median streamflow levels through MAY.

Snow survey measurements show the Salt snowpack to be 62 % of the 30-year average, while combined reservoir storage on the Salt River system is reported at 842,466 acre-feet.



SALT RIVER BASIN
Streamflow Forecasts - February 1, 2004

	<=== Drier === Future Conditions === Wetter ===>						
Forecast Pt	Chance of Exceeding *						
Forecast	90%	70%	50% (Most Prob)	30%	10%	30 Yr Med	
Period	(1000AF)	(1000AF)	(1000AF) (% MED.)	(1000AF)	(1000AF)	(1000AF)	
=====							
Salt River nr Roosevelt							
FEB-MAY	71	132	190	54	262	399	355
FEBRUARY	6.4	15.5	25	54	38	64	46
Tonto Creek ab Gun Creek nr Roosevelt							
FEB-MAY	0.9	6.4	15.0	30	29	63	50
FEBRUARY	1.3	1.5	2.0	16	6.1	16.0	12.6

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average and median are computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.

SALT RIVER BASIN
Reservoir Storage (1000AF) End of January

Reservoir	Usable Capacity	***** This Year	Usable Storage Last Year	***** Average
SALT RIVER RES SYSTEM	2025.8	842.5	558.8	1189.3



SALT RIVER BASIN
Watershed Snowpack Analysis - February 1, 2004

Watershed	Number of Data Sites	This Year as Percent of Last Year	Average
SALT RIVER BASIN	7	161	62

VERDE RIVER BASIN as of February 1, 2004

Well below median streamflow levels are forecast for the basin. In the Verde River, at Horseshoe Dam, the forecast calls for 50 % of median streamflow levels through MAY.

Snow survey measurements show the Verde snowpack to be 50 % of the 30-year average, while combined reservoir storage on the Verde River system is reported to be 117,047 acre-feet.



VERDE RIVER BASIN
Streamflow Forecasts - February 1, 2004

	<=== Drier === Future Conditions === Wetter ===>					
Forecast Pt	Chance of Exceeding *					
Forecast	90%	70%	50% (Most Prob)	30%	10%	30 Yr Med
Period	(1000AF)	(1000AF)	(1000AF) (% MED.)	(1000AF)	(1000AF)	(1000AF)
Verde River abv Horseshoe Dam						
FEB-MAY	36	69	100	50	139	214
FEBRUARY	5.8	15.0	25	71	39	67

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average and median are computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.

VERDE RIVER BASIN
Reservoir Storage (1000AF) End of January

Reservoir	Usable Capacity	***** This Year	***** Usable Storage Last Year	***** Average
VERDE RIVER RES SYSTEM	287.4	117.0	62.7	150.8



VERDE RIVER BASIN
Watershed Snowpack Analysis - February 1, 2004

Watershed	Number of Data Sites	This Year as Percent of Last Year	Percent of Average
VERDE RIVER BASIN	10	202	50
SAN FRANCISCO PEAKS	4	138	69

SAN FRANCISCO-UPPER GILA RIVER BASIN as of February 1, 2004

Well below median streamflow levels are forecast for the basin. In the San Francisco River, at Clifton, the forecast calls for 48 % of median streamflow levels through MAY, while in the Gila River, near Solomon, the forecast calls for 40 % of median streamflow levels through MAY. At San Carlos Reservoir, inflow into the lake is forecast at 36 % of median through MAY.

At San Carlos, reservoir storage stands at 27,264 acre-feet, while snow survey measurements show snowpack levels to be 60 % of the 30-year average.



SAN FRANCISCO - UPPER GILA RIVER BASIN
Streamflow Forecasts - February 1, 2004

	<=== Drier === Future Conditions === Wetter ===>						
Forecast Pt	Chance of Exceeding *						
Forecast	90%	70%	50% (Most Prob)	30%	10%		30 Yr Med
Period	(1000AF)	(1000AF)	(1000AF) (% MED.)	(1000AF)	(1000AF)		(1000AF)
Gila River at Gila							
FEB-MAY	13.6	21	27	51	34	47	53
Gila River nr Virden							
FEB-MAY	7.5	15.0	29	39	51	83	75
San Francisco River at Glenwood							
FEB-MAY	5.0	8.1	10.8	45	14.0	19.9	24
San Francisco River at Clifton							
FEB-MAY	4.7	11.8	28	48	51	84	59
Gila River nr Solomon							
FEB-MAY	14.0	29	57	40	113	194	144
FEBRUARY			16.6	69			24
San Carlos Reservoir inflow							
FEB-MAY	6.7	12.6	30	36	72	134	84

- * 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.
The average and median are computed for the 1971-2000 base period.
- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.

SAN FRANCISCO - UPPER GILA RIVER BASIN
Reservoir Storage (1000AF) End of January

Reservoir	Usable Capacity	***** This Year	***** Usable Storage Last Year	***** Average
SAN CARLOS	875.0	27.2	36.1	421.8
PAINTED ROCK DAM	2492.0	0.0	0.0	166.0



SAN FRANCISCO - UPPER GILA RIVER BASIN
Watershed Snowpack Analysis - February 1, 2004

Watershed	Number of Data Sites	This Year as Percent of Last Year	Average
SAN FRANCISCO - UPPER GILA R	11	164	60

LITTLE COLORADO RIVER BASIN as of February 1, 2004

Well below median streamflow levels are forecast for the basin. In the Little Colorado River, at Lyman Lake, the forecast calls for 39 % of median streamflow levels through JUNE, while at Woodruff, the forecast calls for 39 % of median streamflow levels through MAY.

Additionally, snowpack levels along the southern headwaters of the Little Colorado River, and along the central Mogollon Rim, were measured at 55 % and 57 % of the 30-year average, respectively.



LITTLE COLORADO RIVER BASIN
Streamflow Forecasts - February 1, 2004

	<=== Drier === Future Conditions === Wetter ===>						
Forecast Pt	Chance of Exceeding *						
Forecast	90%	70%	50% (Most Prob)	30%	10%		30 Yr Med
Period	(1000AF)	(1000AF)	(1000AF) (% MED.)	(1000AF)	(1000AF)		(1000AF)
Little Colorado River abv Lyman Lake							
FEB-JUN	0.46	1.53	2.80	39	4.64	8.57	7.10
Rio Nutria nr Ramah							
FEB-MAY	0.04	0.56	1.51	50	3.19	7.42	3.00
Ramah Reservoir inflow							
FEB-MAY	0.08	0.17	0.67	40	2.45	5.05	1.66
Zuni River abv Black Rock Reservoir							
FEB-MAY	0.23	0.45	0.67	49	0.95	1.49	1.36
Little Colorado River at Woodruff							
FEB-MAY	0.28	0.56	1.09	39	2.81	5.29	2.80
Blue Ridge Reservoir inflow							
FEB-MAY	1.3	4.0	6.7	41	10.0	16.2	16.3
Lake Mary inflow							
FEB-MAY	0.44	1.08	1.75	37	2.66	4.49	4.80

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average and median are computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.

(2) - The value is natural volume - actual volume may be affected by upstream water management.

LITTLE COLORADO RIVER BASIN
Reservoir Storage (1000AF) End of January



Reservoir	Usable Capacity	***** This Year	Usable Storage Last Year	***** Average
LYMAN RESERVOIR	30.0	2.1	2.2	14.7
SHOW LOW LAKE	5.1	3.2	2.0	2.9

LITTLE COLORADO RIVER BASIN
Watershed Snowpack Analysis - February 1, 2004

Watershed	Number of Data Sites	This Year as Percent of Last Year	Average
LITTLE COLORADO - SOUTHERN H	8	149	55
CENTRAL MOGOLLON RIM	4	229	57

CHUSKA MOUNTAINS as of February 1, 2004

Snow survey measurements conducted by staffs of the Navajo Tribe show the Chuska snowpack to be 73 % of the long-term average, while well below average streamflow levels are forecast for Captain Tom Wash, Wheatfields Creek, and Bowl Canyon Creek through springtime.



CHUSKA MOUNTAINS
Streamflow Forecasts - February 1, 2004

	<=== Drier === Future Conditions === Wetter ===>						
Forecast Pt	Chance of Exceeding *						
Forecast	90%	70%	50% (Most Prob)	30%	10%		30 Yr Avg
Period	(1000AF)	(1000AF)	(1000AF) (% AVG.)	(1000AF)	(1000AF)		(1000AF)
Captain Tom Wash nr Two Gray Hills							
MAR-MAY	0.42	0.85	1.70	60	3.40	5.90	2.83
Wheatfields Creek nr Wheatfields							
MAR-MAY	0.44	0.87	1.75	60	3.55	6.05	2.90
Bowl Canyon Creek abv Assayi Lake							
MAR-MAY	0.15	0.30	0.60	60	1.20	2.05	1.00

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.



CHUSKA MOUNTAINS
Watershed Snowpack Analysis - February 1, 2004

Watershed	Number of Data Sites	This Year as Percent of Last Year	Average
CHUSKA MOUNTAINS	7	135	73
DEFIANCE PLATEAU	2	422	67

NORTHWESTERN ARIZONA as of February 1, 2004

On the Colorado River, inflow into Lake Powell is forecast at 82 % of the 30-year average through JULY, while at Littlefield, the Virgin River is forecast at 47 % of average through JULY.

At the Grand Canyon, snow survey measurements conducted by the National Park Service show the snowpack to be at 73 % of the 30-year average.



NORTHWESTERN ARIZONA
Streamflow Forecasts - February 1, 2004

	<=== Drier === Future Conditions === Wetter ===>						
Forecast Pt	Chance of Exceeding *						
Forecast	90%	70%	50% (Most Prob)	30%	10%		30 Yr Avg
Period	(1000AF)	(1000AF)	(1000AF) (% AVG.)	(1000AF)	(1000AF)		(1000AF)
Virgin River at Littlefield							
APR-JUL	12.9	26	35	47	44	57	74
Lake Powell inflow							
APR-JUL	3420	5260	6500	82	7740	9580	7930

* 90%, 70%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

- (1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
- (2) - The value is natural volume - actual volume may be affected by upstream water management.

NORTHWESTERN ARIZONA
Reservoir Storage (1000AF) End of January

Reservoir	Usable Capacity	***** This Year	Usable Storage Last Year	***** Average
LAKE HAVASU	619.0	510.6	537.1	551.8
LAKE MOHAVE	1810.0	1623.4	1705.3	1672.3
LAKE MEAD	26159.0	15434.0	16854.0	21992.0
LAKE POWELL	24322.0	10984.0	13269.0	18463.0

NORTHWESTERN ARIZONA
Watershed Snowpack Analysis - February 1, 2004

Watershed	Number of Data Sites	This Year as Percent of Last Year	Average
GRAND CANYON	1	190	73

S N O W S U R V E Y D A T A

FEBRUARY 1, 2004

SNOW COURSE	ELEV.	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 71-00
ARBABS FOREST (AK)	7680	1/29	6	1.6	0.9	2.6
BAKER BUTTE SNOTEL	7330	2/01	-	2.5	0.1	4.6
BAKER BUTTE #2	7700	1/30	14	3.8	2.8	8.2
BALDY SNOTEL	9220	2/01	-	3.5	4.1	5.7
BEAR PAW	10100	1/29	31	9.7	6.4	12.2
BEAVER HEAD	8000	1/30	6	1.4	0.0	2.8
BEAVER HEAD SNOTEL	7990	2/01	-	1.5	2.4	3.1
BEAVER SPRING	9220	1/28	20	4.5	3.2	7.5
BRIGHT ANGEL	8400	1/30	18	5.5	2.9	7.5
BUCK SPRING	7400	2/05	11	1.4	0.0	4.8
CHALENDER	7100	1/28	1	0.3	0.0	2.5
CHEESE SPRINGS	8600	1/28	13	2.4	2.9	4.3
CORONADO TRL SNOTEL	8400	2/01	-	1.7	0.0	3.2
CORONADO TRAIL	8350	1/30	4	1.1	0.0	2.6
FLUTED ROCK	7800	1/29	10	2.2	0.0	3.1
FORT APACHE	9160	1/28	20	3.6	3.9	6.1
FORT VALLEY	7350	1/28	2	0.7	0.0	2.4
FRY SNOTEL	7220	2/01	-	2.9	2.5	4.9
GRAND CANYON	7500	no report			1.3	2.6
HANNAGAN MDWS SNOTEL	9020	2/01	-	5.8	3.5	8.6
HAPPY JACK	7630	1/29	6	1.2	0.0	3.8
HAPPY JACK SNOTEL	7630	2/01	-	3.0	2.7	3.7
HEBER SNOTEL	7640	2/01	-	3.5	0.0	4.8
LAKE MARY	6930	1/30	6	1.3	0.0	2.7
MAVERICK FORK SNOTEL	9200	2/01	-	4.3	4.5	7.3
MORMON MTN SNOTEL	7500	2/01	-	3.0	0.0	4.9
MORMON MT. SUMMIT #2	8470	1/30	20	4.9	3.7	9.0
NEWMAN PARK	6750	1/28	5	1.1	0.0	2.5
NUTRIOSO	8500	1/30	4	1.4	0.0	1.7
PROMONTORY SNOTEL	7900	2/01	-	5.8	3.9	9.7
SNOW BOWL #1 ALT.	10260	1/28	17	3.0	1.3	8.7
SNOW BOWL #2	11000	1/28	27	6.4	3.6	11.8
SNOWSLIDE CANYON	9750	1/29	26	7.2	6.1	9.5
SNOWSLIDE CYN SNOTEL	9750	2/01	-	9.6	9.5	9.1
TSAILE CANYON #1	8160	1/27	18	4.0	3.0	5.3
TSAILE CANYON #3	8920	1/27	26	6.1	5.2	7.2
WHITE HORSE SNOTEL	7180	2/01	-	0.8	0.9	3.8
WILDCAT SNOTEL	7850	2/01	-	1.8	0.0	3.4
WILLIAMS SKI RUN	7720	1/28	12	3.0	1.1	6.1
WORKMAN CREEK SNOTEL	6900	2/01	-	5.7	0.9	4.8